



**Brighter strategies**  
for greener projects



Client: Lovell Homes  
Project: Novartis Phase 1 & 2  
Report: Invertebrate Survey Report

## QUALITY ASSURANCE

Issue/Revision:	Draft	Final
Date:	October 2025	October 2025
Comments:		
Prepared by:	Jess Cole	Jess Cole
Authorised by:	Mike Harris	Mitch Cooke
File Reference:	552979jc17Jul25FV01_Invertebrates	552979jc17Jul25FV01_Invertebrates

## CONTENTS

<b>1.0 EXECUTIVE SUMMARY</b>	<b>1</b>
<b>2.0 INTRODUCTION</b>	<b>2</b>
2.1 SITE DESCRIPTION	2
2.2 ECOLOGICAL BACKGROUND	3
<b>3.0 METHODOLOGY</b>	<b>4</b>
3.1 COMPETENCIES	5
3.2 CONSTRAINTS	6
<b>4.0 RESULTS</b>	<b>7</b>
4.1 SURVEY 1 - 23RD MAY 2025	7
4.2 SURVEY 2 - 6TH AUGUST 2025	7
<b>5.0 DISCUSSION &amp; RECCOMENDATION</b>	<b>8</b>
5.1 ENHANCEMENT	8
<b>6.0 SUMMARY &amp; CONCLUSIONS</b>	<b>10</b>
<b>APPENDIX A FULL SPECIES LIST</b>	
<b>REFERENCES</b>	

### Tables

Table A.1	Full speices list - May 2025
Table A.2	Full species list - August 2025

### Figures

No table of figures entries found.

### Plates

Plate 3.1	Novartis Phase 1 &2 and Novartis Phase 3	5
Plate 5.1	Example of loggery	8
Plate 5.2	Example of habitat panels	9
Plate 5.3	Example of pollinator posts	9

## 1.0 EXECUTIVE SUMMARY

Greengage Environmental (Ltd) were commissioned by Lovell Homes to undertake an invertebrate survey at the site known as Novartis Phase 1 & 2, Horsham, West Sussex, hereafter referred to as 'the site'.

The survey work was commissioned to inform the planning application for the site (Planning Ref: DC/25/0415) which seeks the retention and change of use of the current building on-site, and erection of multiple resident units (with associated infrastructure).

Greengage completed a Preliminary Ecological Appraisal (PEA)<sup>1</sup> of the site on 5th November 2024 and 12th February 2025, which included a site survey and analysis of records from Sussex Biodiversity Records Centre (SxBRC) within 2km of the site. Through the site survey, high suitability for the site to support common and widespread and potentially notable invertebrates, owing to the number of local records in the areas and mosaic of habitats present, was recorded. Full assessment of the site's suitability for invertebrates and the desk study records can be found within the PEA report<sup>1</sup>.

It was initially recommended that a suite of three surveys by a specialist entomologist should be carried out across the site between May and July to confirm the presence/likely absence of any nationally scarce invertebrates on site. However, after discussion with an experienced entomologist, it was decided that a single baseline survey should first be conducted and the need for any further survey determined thereafter. Following this baseline survey, based on the advice of the suitable experienced entomologist, a second survey was then conducted to reinforce the results collected during the first survey.

This document is a report of the survey results and provides discussion around best practice enhancement measures for invertebrates on site post development.

A total of 97 invertebrate taxa were identified during the first survey and 226 during the second. Species recorded were mostly common though some notable species were also recorded, including Section 41 Priority Species and Red Data Book (RDB)<sup>2</sup> species.

Based on the above results mitigation measures for invertebrates are not considered necessary. Nevertheless, measures to compensate for the loss of habitat on site should be provided, this should be provided through the creation of areas of species rich grasslands (meadows) maintained with areas of both long and short sward heights and areas of bare ground, as well as provision of areas of scrub.

Enhancement measures for the benefit of invertebrates are also recommended for implementation post development., including provision of loggeries, habitat panels and pollinator posts. A Landscape Enhancement Management Plan (LEMP) providing greater detail on these compensation and enhancement features in accordance with British Standard 42020: 2013 Biodiversity<sup>3</sup>, should be produced as part of a planning condition.

The data collected during the bat emergence survey is considered valid for 12-18 months in accordance Chartered Institute of Ecology and Environmental Management (CIEEM) guidance<sup>4</sup> and an updated survey will be required if the works have not started within this timeframe or should site conditions change significantly during this time.

## 2.0 INTRODUCTION

Greengage were commissioned by Lovell Homes to undertake an invertebrate survey at the site known as Novartis Phase 1 & 2, Horsham, West Sussex, hereafter referred to as 'the site'.

The survey work was commissioned to inform the planning application for the site (Planning Ref: DC/25/0415) which seeks the retention and change of use of the current building on-site, and erection of multiple resident units (with associated infrastructure).

This document is a report of the survey results and provides discussion around best practice enhancement measures for invertebrates on site post development.

### 2.1 SITE DESCRIPTION

The site extends to approximately 2.63 hectares (ha) and is centred on National Grid Reference TQ 17809 31816, OS Co-ordinates 517809, 131816.

The site is comprised primarily of developed land; sealed surface, with one large existing building located adjacent to the site's eastern boundary which contains a courtyard hereafter referred to as "the former Novartis building". The courtyard included a pond, surrounded by bramble scrub, dense scrub, modified grassland and individual trees. Two large patches of sparsely vegetated urban land were located towards the centre of the site, both of which were bordered by large patches of bramble scrub. Multiple areas of other neutral grassland were located throughout the site, positioned around the centre of the site, and in the northeast and southwest corners of the site. Two patches of willow scrub were located adjacent to the western site boundary, with an area of modified grassland located towards the northwest corner. At the entrance to the site along the western boundary, a small patch of other woodland - mixed - mainly conifer was present to the south, with bramble scrub to the north bordered by another native hedgerow. Individual trees were located throughout the site, with the highest density located along the northern boundary of the site.

The site is located in the centre of Horsham and therefore situated in an urban setting, primarily surrounded by residential buildings and gardens. Parsonage road and Wimblehurst road run along the northern and western boundaries of the site respectively, with a railway line running adjacent to the southern boundary, with an additional railway line located in close proximity to the east of the site.

Fragmented priority woodland is found throughout Horsham with the closest found in Horsham Park approximately 480 metres (m) south of the site boundary. Warnham Local Nature Reserve (LNR) is located approximately 665 m northwest of the site boundary, with a golf course located directly south of the LNR. Large areas of ancient woodland can be found within the wider area, with the closest located in approximately 850 m north of the site boundary. Multiple parcels of different priority habitats are located between 1 km to 2 km from the site boundary. These include woodland pasture and parks, good quality semi-improved grassland (non priority), ancient replacement woodland, and lowland meadows which are all classified as priority habitats.

---

## 2.2 ECOLOGICAL BACKGROUND

### 2018-2022

An initial Ecological Appraisal by Hampshire County Council originally produced in 2015 and updated in 2018/2019<sup>5</sup> surveyed across the whole Novartis site, incorporating what is now known as Novartis Phase 1&2 and Novartis Phase 3.

A subsequent PEA was then produced by Ecology & Habitat Management Ltd<sup>6</sup> in 2022 which assessed the site. The Phase 1 habitats survey undertaken as part of the PEA identified the habitats of building, hardstanding, bare ground, semi-improved grassland, scrub, introduced shrub, and scattered tree across the site.

Both of these surveys recognised value for invertebrates though no further survey for invertebrates was conducted.

### 2024

In November 2024 Greengage completed an updated PEA<sup>1</sup> of the site. which identified high suitability for the site to support invertebrates, including notable invertebrates, due to the types of habitats present on site and the records returned from Sussex Biodiversity Records Centre (SxBRC). Off the back of this, it was initially recommended that a suite of three surveys by a specialist entomologist should be carried out across the site between May and July to confirm the presence/likely absence of any nationally scarce invertebrates on site. However, after discussion with an experienced entomologist, it was decided that a single baseline survey should first be conducted and the need for any further survey determined thereafter.

### 3.0 METHODOLOGY

The baseline invertebrate survey was completed on 23rd May 2025. The second survey was completed on 6th August 2025. The surveys were conducted in appropriate weather conditions with temperatures above 15°C and no rain.

Due to it being impracticable to survey all the potential invertebrates within any given site, only specific groups of species were examined during fieldwork. These groups are sufficiently well known as to allow meaningful comparisons to be made with other sites, both locally and nationally. They are also important as indicators of the quality of a site and the habitats present<sup>7</sup>.

Groups covered during the survey were:

- *Mollusca* (slugs and snails)
- *Arachnida* (spiders, harvestmen & pseudoscorpions)
- *Isopoda* (woodlice)
- *Thysanura* (bristletails)
- *Odonata* (dragonflies & damselflies)
- *Orthoptera* (grasshoppers & crickets)
- *Dictyoptera* (cockroaches)
- *Dermaptera* (earwigs)
- *Hemiptera-Heteroptera* (true-bugs)
- *Hemiptera-Homoptera* (hoppers)
- *Neuroptera* (lace-wings)
- *Mecoptera* (scorpion-flies)
- *Lepidoptera* (butterflies & moths)
- *Diptera* (true flies)
- *Aculeate Hymenoptera* (ants, bees & wasps)
- *Coleoptera* (beetles)

Standard field techniques were employed to sample the terrestrial and arboreal invertebrates, namely sweep netting, beating foliage over a beating tray etc. In addition, a battery powered suction sampler was employed to sample terrestrial invertebrates.

N.B it should be noted that the results presented below are species recorded across both the Novartis Phase 1 & 2 site and the neighbouring Novartis Phase 3 site. The locations of both plots are shown in Plate 3.1 below. Data is presented this way as both plots have similar habitat types and there is barrier to movement of inverts between the two sites.

Plate 3.1 Novartis Phase 1 &amp; 2 and Novartis Phase 3



### 3.1 COMPETENCIES

The baseline invertebrate survey was completed on 23rd May 2025 by experienced entomologist Dr Jonty Denton. Dr Denton is a freelance Chartered Ecologist with over 30 years experience, with Natural England licenses for Bats, Dormice, Great Crested Newt, Natterjacks, Sand Lizard, Smooth Snake, and White-clawed Crayfish. His clients include Natural England, the National Trust, Crown Estates, County Trusts, Butterfly Conservation, the Ministry of Defence, Royal Parks and many County and District Councils, as well as the Environment Agency and Thames Water.

The second survey was completed on 6th August 2025 conducted by Martin Townsend BSc, Zoology (Aberdeen). Martin was elected Fellow of the Royal Entomological Society and holds full Membership to of the Chartered Institute of Ecology and Environmental Management (CIEEM), British Entomological and Natural History Society, Amateur Entomologists' Society and British Ecological Society. Martin has over 20 years' experience in ecological consultancy, carrying out and managing general and targeted invertebrate surveys in the private and public sectors, for planning, conservation management, control of Invasive Non-native Species (INNS) and ecological research. Martin also has over 25 years in ecological research, conservation and scientific publishing. Recognised authority and published author of identification guides on Lepidoptera.

Jess Cole, Senior Consultant, who wrote this report, has a BSc degree in Ecology (Hons) and is an Associate member of CIEEM. Jess holds a Natural England Great Crested Newt Licence and has over seven years' experience in ecological survey and assessment.

Mike Harris, Director, who reviewed this report, has a Bachelor's degree in Environmental Biology (BSc Hons), a Natural England Great Crested Newt Licence and Dormouse Licence, is a Chartered Environmentalist (CEnv) and Full member of CIEEM. Mike has over 20 years' experience in ecological surveying and has undertaken and managed numerous ecological surveys and assessments.

Mitch Cooke, Director, who verified this report, has a degree in Ecology (Hons), an MSc in Environmental Assessment and Management, and is a Full member of CIEEM with over 35 years' experience in ecological survey and assessment. Mitch has set up and developed ecological and environmental teams for nearly 20 years and has undertaken and managed numerous ecological surveys and assessments. He is the Director at Greengage and manages the team.

This report was written by Jess Cole, reviewed by Francesca Thorley and verified by Mitch Cooke who confirms in writing (see the QA sheet at the front of this report) that the report is in line with the following:

- Represents sound industry practice;
- Reports and recommends correctly, truthfully and objectively;
- Is appropriate given the local site conditions and scope of works proposed; and
- Avoids invalid, biased and exaggerated statements.

## 3.2 CONSTRAINTS

The surveys were conducted at a suitable time of year by suitably qualified entomologists. All of the site was accessible at the time of the surveys.

## 4.0 RESULTS

A full species list from both surveys can be found within Appendix A of this letter.

### 4.1 SURVEY 1 - 23RD MAY 2025

A total of 97 invertebrate taxa were identified during this survey.

Of these 97, two of the species have a nationally significant status. This is not considered to be a high number for a brownfield site in southern England.

The two species identified with nationally significant status included cinnabar moth *Tyria jacobaeae* and small heath *Coenonympha pamphilus*. These are both Section 41 Priority Species, however cinnabar is considered to be a 'research only' species only, meaning the species has the potential to be under threat owing to declines or some other reason such as eradication of their foodplant (e.g. cinnabar moth and its host plant, ragwort) but not sufficiently scarce or under threat enough yet to be classed as scarce.

Five other species recorded on site are classified as Nationally Scarce Category B - Notable B (Nb) or Nationally Scarce (NS), but do not fall within the Red Data Book (RDB)<sup>2</sup> these were *Diplapion stolidum*, *Magdalis cerasi*, *Orthochaetes setiger*, *Drilus flavescens* and *Bathysolen nubilus*.

The remaining 90 species were either nationally or locally common.

### 4.2 SURVEY 2 - 6TH AUGUST 2025

During the August survey a total of 226 species were recorded, of these 26 were recorded previously during the May survey, including cinnabar moth and small heath.

Of the 200 newly recorded species 12 were notable in some way. Eight were Nb or NS including Dusky Cockroach *Ectobius lapponicus*, *Rhopalus parumpunctatus*, *Anoscopus albiger*, *Protapion difforme*, *Protapion filirostre*, *Licinus punctatulus*, *Glocianus punctiger* and *Trichosirocalus barnevillei*. One species, *Terellia longicauda*, was classified as scarce and *Orchisia costata* and *Blaesoxipha plumicornis* was potentially NS (pNS) and potentially near threatened (pNT). Lastly. One species, *Nysius graminicola*, was listed within category 3 of the RDB<sup>2</sup>. Category 3 species are taxa with small populations in Great Britain that are not at present endangered or vulnerable, but are at risk.

All other 188 species recorded in August, which were not previously, recorded in May, were nationally common.

## 5.0 DISCUSSION & RECCOMENDATION

No further survey is considered necessary, in light of the data collected

More than twice as many species are associated with tall sward grassland and scrub compared to species associated with short sward grassland and bare ground. However, analysing species with elevated status, roughly twice as many are associated with short sward grassland and bare ground species as are associated with tall sward grassland and scrub. This pattern is not untypical of a site with these mosaics of habitat.

Based on the above results mitigation measures for invertebrates are not considered necessary. Nevertheless, measures to compensate for the loss of habitat on site should be provided, this should be provided though the creation of area of species rich grasslands (meadows) maintained with areas of both long and short sward heights and areas of bare ground, as well as provision of areas of scrub.

### 5.1 ENHANCEMENT

There is also opportunity to enhance the site for invertebrates through the provision of the following features. A Landscape Enhancement Management Plan (LEMP) providing greater detail on the above compensation and enhancement features in accordance with British Standard 42020: 2013 Biodiversity<sup>3</sup>, should be produced and implemented for the site and secured through planning condition.

#### Loggeries

Loggeries should be created within the areas of planting. Log sizes should range from ~10cm up to ~40cm diameter with approximately one third of the logs buried. Plants such as ferns, bulbs, and other woodland understorey plants should then be planted amongst the loggeries.

Wood from broadleaved trees such as oak and beech, and from fruiting trees such as apple and pear where possible should be used, at least 100mm in diameter with the bark still on. Logs in contact with the substrate should remain damp underneath, which is vital for many invertebrates such as woodlice. Logs should be placed both vertically and horizontally in clusters; vertical standing wood should be incorporated by submerging the logs into the full depth of the substrate.

Plate 5.1 Example of loggery



## Habitat Panels

Habitat panels should be located amongst areas of nectar rich planting or on areas of biodiverse green roof, so to provide a constant nectar resource for pollinators on a year-round basis. Habitat panels should be included in sunny, south-facing areas.

Panels should use untreated wood products which provide a range of opportunities for sheltering and nesting solitary bees and other invertebrates. They can be 'designed' or integrated with interpretation panels. Design of each panel should avoid providing opportunities for numerous taxa to minimise risk of parasitism.

Plate 5.2 Example of habitat panels



## Pollinator posts

Similar to habitat panels, pollinator posts should be located amongst areas of nectar rich planting or installed on extensive biodiverse green roof. These should be close to planting in order to provide a constant nectar resource for pollinators on a year-round basis and located in sunny, south-facing areas within landscaped areas and public realm.

Plate 5.3 Example of pollinator posts



## 6.0 SUMMARY & CONCLUSIONS

Greengage were commissioned by Lovell Homes to undertake an invertebrate survey at the site known as Novartis Phase 1 & 2, Horsham, West Sussex.

Greengage completed a PEA<sup>1</sup> of the site on 5th November 2024 and 12th February 2025, which included a site survey and analysis of records from SxBRC within 2km of the site. Through the site survey, high suitability for the site to support common and widespread and potentially notable invertebrates, owing to the number of local records in the areas and mosaic of habitats present, was recorded.

It was initially recommended that a suite of three surveys by a specialist entomologist should be carried out across the site between May and July to confirm the presence/likely absence of any nationally scarce invertebrates on site. However, after discussion with an experienced entomologist, it was decided that a single baseline survey should first be conducted and the need for any further survey determined thereafter. Following this baseline survey, based on the advice of the suitable experienced entomologist, a second survey was then conducted to reinforce the results collected during the first survey.

A total of 97 invertebrate taxa were identified during the first survey and 226 during the second. Species recorded were mostly common though some notable species were also recorded, including Section 41 Priority Species and RDB<sup>2</sup> species.

Based on the above results mitigation measures for invertebrates are not considered necessary. Nevertheless, measures to compensate for the loss of habitat on site should be provided, this should be provided through the creation of areas of species rich grasslands (meadows) maintained with areas of both long and short sward heights and areas of bare ground, as well as provision of areas of scrub.

Section 5 of this report also recommends enhancement measures for the benefit of invertebrates post development.

A LEMP providing greater detail on these compensation and enhancement features in accordance with British Standard 42020: 2013 Biodiversity<sup>3</sup>, should be produced as part of a planning condition.

The data collected during the bat emergence survey is considered valid for 12-18 months in accordance Chartered Institute of Ecology and Environmental Management (CIEEM) guidance<sup>4</sup> and an updated survey will be required if the works have not started within this timeframe or should site conditions change significantly during this time.

## APPENDIX A FULL SPECIES LIST

Table A.1 Full speices list - May 2025

Species	Family	Order
<i>Eratigena duellica</i>	Agelenidae	Araneae
<i>Anyphaena accentuata</i>	Anyphaenidae	Araneae
<i>Araneus diadematus</i>	Araneidae	Araneae
<i>Mangora acalypha</i>	Araneidae	Araneae
<i>Zygiella x-notata</i>	Araneidae	Araneae
<i>Brigittea latens</i>	Dictynidae	Araneae
<i>Dysdera crocata</i>	Dysderidae	Araneae
<i>Pocadicnemis juncea</i>	Linyphiidae	Araneae
<i>Philodromus albidus</i>	Philodromidae	Araneae
<i>Philodromus rufus sens. str.</i>	Philodromidae	Araneae
<i>Pisaura mirabilis</i>	Pisauridae	Araneae
<i>Euophrys frontalis</i>	Salticidae	Araneae
<i>Heliophanus cupreus</i>	Salticidae	Araneae
<i>Heliophanus flavipes</i>	Salticidae	Araneae
<i>Salticus scenicus</i>	Salticidae	Araneae
<i>Diplapion stolidum</i>	Apionidae	Coleoptera
<i>Exapion ulicis</i>	Apionidae	Coleoptera
<i>Ischnopteration loti</i>	Apionidae	Coleoptera
<i>Malthodes marginatus</i>	Cantharidae	Coleoptera
<i>Bembidion lampros</i>	Carabidae	Coleoptera
<i>Syntomus foveatus</i>	Carabidae	Coleoptera
<i>Batophila aerata</i>	Chrysomelidae	Coleoptera
<i>Bruchidius cisti</i>	Chrysomelidae	Coleoptera
<i>Cryptocephalus moraei</i>	Chrysomelidae	Coleoptera
<i>Plagiodera versicolora</i>	Chrysomelidae	Coleoptera
<i>Coccinella septempunctata</i>	Coccinellidae	Coleoptera
<i>Harmonia axyridis</i>	Coccinellidae	Coleoptera
<i>Psyllobora vigintiduopunctata</i>	Coccinellidae	Coleoptera
<i>Magdalis cerasi</i>	Curculionidae	Coleoptera
<i>Mecinus pascuorum</i>	Curculionidae	Coleoptera
<i>Orthochaetes setiger</i>	Curculionidae	Coleoptera

Species	Family	Order
<i>Sitona striatellus</i>	Curculionidae	Coleoptera
<i>Tychius junceus</i>	Curculionidae	Coleoptera
<i>Drilus flavescens</i>	Drilidae	Coleoptera
<i>Cordylepherus viridis</i>	Malachiidae	Coleoptera
<i>Malachius bipustulatus</i>	Malachiidae	Coleoptera
<i>Meligethes carinulatus</i>	Nitidulidae	Coleoptera
<i>Oedemera lurida</i>	Oedemeridae	Coleoptera
<i>Oedemera nobilis</i>	Oedemeridae	Coleoptera
<i>Ochina ptinoides</i>	Ptinidae	Coleoptera
<i>Contacyphon variabilis</i>	Scirtidae	Coleoptera
<i>Anaspis maculata</i>	Scraptiidae	Coleoptera
<i>Anaspis regimbarti</i>	Scraptiidae	Coleoptera
<i>Drusilla canaliculata</i>	Staphylinidae	Coleoptera
<i>Stenus aceris</i>	Staphylinidae	Coleoptera
<i>Stenus clavicornis</i>	Staphylinidae	Coleoptera
<i>Tasgius melanarius</i>	Staphylinidae	Coleoptera
<i>Leptogaster cylindrica</i>	Asilidae	Diptera
<i>Lucilia sericata</i>	Calliphoridae	Diptera
<i>Chrysotoxum festivum</i>	Syrphidae	Diptera
<i>Eristalis pertinax</i>	Syrphidae	Diptera
<i>Merodon equestris</i>	Syrphidae	Diptera
<i>Urophora stylata</i>	Tephritidae	Diptera
<i>Cardiastethus fasciiventris</i>	Anthocoridae	Hemiptera
<i>Acericerus ribauti</i>	Cicadellidae	Hemiptera
<i>Bathysolen nubilus</i>	Coreidae	Hemiptera
<i>Tachycixius pilosus</i>	Cixiidae	Hemiptera
<i>Kleidocerys resedae</i>	Lygaeidae	Hemiptera
<i>Nysius huttoni</i>	Lygaeidae	Hemiptera
<i>Scolopostethus decoratus</i>	Lygaeidae	Hemiptera
<i>Taphropeltus contractus</i>	Lygaeidae	Hemiptera
<i>Centrotus cornutus</i>	Membracidae	Hemiptera
<i>Capsus ater</i>	Miridae	Hemiptera
<i>Closterotomus trivialis</i>	Miridae	Hemiptera
<i>Deraeocoris lutescens</i>	Miridae	Hemiptera

Species	Family	Order
<i>Phylus melanocephalus</i>	Miridae	Hemiptera
<i>Pinalitus cervinus</i>	Miridae	Hemiptera
<i>Psallus perresi</i>	Miridae	Hemiptera
<i>Psallus varians</i>	Miridae	Hemiptera
<i>Palomena prasina</i>	Pentatomidae	Hemiptera
<i>Pentatoma rufipes</i>	Pentatomidae	Hemiptera
<i>Kalama tricornis</i>	Tingidae	Hemiptera
<i>Lasius niger</i>	Formicidae	Hymenoptera
<i>Myrmica ruginodis</i>	Formicidae	Hymenoptera
<i>Amblyteles armatorius</i>	Ichneumonidae	Hymenoptera
<i>Ammophila sabulosa</i>	Sphecidae	Hymenoptera
<i>Rhogogaster viridis</i> (after Taeger & Viitasaari, 2015)	Tenthredinidae	Hymenoptera
<i>Tenthredo livida</i>	Tenthredinidae	Hymenoptera
<i>Armadillidium vulgare</i>	Armadillidiidae	Isopoda
<i>Euclidia glyphica</i>	Erebidae	Lepidoptera
<i>Orgyia antiqua</i>	Erebidae	Lepidoptera
<i>Tyria jacobaeae</i>	Erebidae	Lepidoptera
<i>Cameraria ohridella</i>	Gracillariidae	Lepidoptera
<i>Polyommatus icarus</i>	Lycaenidae	Lepidoptera
<i>Coenonympha pamphilus</i>	Nymphalidae	Lepidoptera
<i>Pieris rapae</i>	Pieridae	Lepidoptera
<i>Dichrorampha alpinana</i>	Tortricidae	Lepidoptera
<i>Zygaena filipendulae</i>	Zygaenidae	Lepidoptera
<i>Libellula depressa</i>	Libellulidae	Odonata
<i>Valenzuela flavidus</i>	Caeciliusidae	Psocodea
<i>Graphopsocus cruciatus</i>	Stenopsocidae	Psocoptera
<i>Arion (Arion) rufus</i>	Arionidae	Pulmonata
<i>Cepaea (Cepaea) nemoralis</i>	Helicidae	Pulmonata
<i>Cornu aspersum</i>	Helicidae	Pulmonata
<i>Hygromia (Hygromia) cinctella</i>	Hygromiidae	Pulmonata
<i>Limacus flavus</i>	Limacidae	Pulmonata
<i>Pupilla (Pupilla) muscorum</i>	Pupillidae	Pulmonata

Table A.2 Full species list - August 2025

Species	Family	Order
<i>Agalena labyrinthica</i>	Agelenidae	Araneae
<i>Agalenatea redii</i>	Araneidae	Araneae
<i>Araneus diadematus</i>	Araneidae	Araneae
<i>Argiope bruennichi</i>	Araneidae	Araneae
<i>Nuctenea umbratica</i>	Araneidae	Araneae
<i>Clubiona corticalis</i>	Clubionidae	Araneae
<i>Dysdera crocata</i>	Dysderidae	Araneae
<i>Micaria micans</i>	Gnaphosidae	Araneae
<i>Zelotes latreillei</i>	Gnaphosidae	Araneae
<i>Stemonyphantes lineatus</i>	Linyphiidae	Araneae
<i>Tenuiphantes tenuis</i>	Linyphiidae	Araneae
<i>Pardosa pullata</i>	Lycosidae	Araneae
<i>Pisaura mirabilis</i>	Pisauridae	Araneae
<i>Heliophanus cupreus</i>	Salticidae	Araneae
<i>Heliophanus flavipes</i>	Salticidae	Araneae
<i>Pachygnatha degeeri</i>	Tetragnathidae	Araneae
<i>Enoplognatha latimana</i>	Theridiidae	Araneae
<i>Enoplognatha ovata sens. str.</i>	Theridiidae	Araneae
<i>Ozyptila simplex</i>	Thomisidae	Araneae
<i>Chorthippus parallelus</i>	Acrididae	Orthoptera
<i>Chorthippus albomarginatus</i>	Acrididae	Orthoptera
<i>Chorthippus brunneus</i>	Acrididae	Orthoptera
<i>Omocestus viridulus</i>	Acrididae	Orthoptera
<i>Ectobius lapponicus</i>	Blattellidae	Blattodea
<i>Conocephalus fuscus</i>	Tettigoniidae	Orthoptera
<i>Anthocoris nemorum</i>	Anthocoridae	Hemiptera
<i>Orius majusculus</i>	Anthocoridae	Hemiptera
<i>Berytinus minor</i>	Berytidae	Hemiptera
<i>Coreus marginatus</i>	Coreidae	Hemiptera
<i>Cymus melanocephalus</i>	Lygaeidae	Hemiptera
<i>Kleidocerys resedae</i>	Lygaeidae	Hemiptera
<i>Nysius graminicola</i>	Lygaeidae	Hemiptera

Species	Family	Order
<i>Nysius huttoni</i>	Lygaeidae	Hemiptera
<i>Nysius senecionis</i>	Lygaeidae	Hemiptera
<i>Stygnocoris rusticus</i>	Lygaeidae	Hemiptera
<i>Stygnocoris sabulosus</i>	Lygaeidae	Hemiptera
<i>Taphropeltus contractus</i>	Lygaeidae	Hemiptera
<i>Blepharidopterus angulatus</i>	Miridae	Hemiptera
<i>Deraeocoris lutescens</i>	Miridae	Hemiptera
<i>Megacoelum infusum</i>	Miridae	Hemiptera
<i>Orthops campestris</i>	Miridae	Hemiptera
<i>Phytocoris varipes</i>	Miridae	Hemiptera
<i>Stenodema calcarata</i>	Miridae	Hemiptera
<i>Stenodema laevigata</i>	Miridae	Hemiptera
<i>Himacerus mirmicoides</i>	Nabidae	Hemiptera
<i>Nabis limbatus</i>	Nabidae	Hemiptera
<i>Dolycoris baccarum</i>	Pentatomidae	Hemiptera
<i>Palomena prasina</i>	Pentatomidae	Hemiptera
<i>Pentatoma rufipes</i>	Pentatomidae	Hemiptera
<i>Piezodorus lituratus</i>	Pentatomidae	Hemiptera
<i>Corizus hyoscyami</i>	Rhopalidae	Hemiptera
<i>Myrmus miriformis</i>	Rhopalidae	Hemiptera
<i>Rhopalus parumpunctatus</i>	Rhopalidae	Hemiptera
<i>Rhopalus subrufus</i>	Rhopalidae	Hemiptera
<i>Stictopleurus punctatonervosus</i>	Rhopalidae	Hemiptera
<i>Eurygaster testudinaria</i>	Scutelleridae	Hemiptera
<i>Tingis cardui</i>	Tingidae	Hemiptera
<i>Philaenus spumarius</i>	Aphrophoridae	Hemiptera
<i>Aphrophora alni</i>	Aphrophoridae	Hemiptera
<i>Anoscopus albifrons</i>	Cicadellidae	Hemiptera
<i>Anoscopus albiger</i>	Cicadellidae	Hemiptera
<i>Aphrodes diminuta</i>	Cicadellidae	Hemiptera
<i>Athysanus argentarius</i>	Cicadellidae	Hemiptera
<i>Cicadella viridis</i>	Cicadellidae	Hemiptera
<i>Conosanus obsoletus</i>	Cicadellidae	Hemiptera
<i>Eupelix cuspidata</i>	Cicadellidae	Hemiptera

Species	Family	Order
<i>Eupteryx aurata</i>	Cicadellidae	Hemiptera
<i>Eupteryx melissae</i>	Cicadellidae	Hemiptera
<i>Euscelis incisus</i>	Cicadellidae	Hemiptera
<i>Iassus lanio</i>	Cicadellidae	Hemiptera
<i>Macropsis scotti</i>	Cicadellidae	Hemiptera
<i>Macrosteles sexnotatus</i>	Cicadellidae	Hemiptera
<i>Megophthalmus scanicus</i>	Cicadellidae	Hemiptera
<i>Mocydia crocea</i>	Cicadellidae	Hemiptera
<i>Oncopsis appendiculata</i>	Cicadellidae	Hemiptera
<i>Populicerus confusus</i>	Cicadellidae	Hemiptera
<i>Psammotettix confinis</i>	Cicadellidae	Hemiptera
<i>Streptanus sordidus</i>	Cicadellidae	Hemiptera
<i>Zyginidia scutellaris</i>	Cicadellidae	Hemiptera
<i>Conomelus anceps</i>	Delphacidae	Hemiptera
<i>Dicranotropis hamata</i>	Delphacidae	Hemiptera
<i>Javesella pellucida</i>	Delphacidae	Hemiptera
<i>Aspidapion aeneum</i>	Apionidae	Coleoptera
<i>Eutrichapion vorax</i>	Apionidae	Coleoptera
<i>Ischnopterapion loti</i>	Apionidae	Coleoptera
<i>Oxystoma pomonae</i>	Apionidae	Coleoptera
<i>Protapion difforme</i>	Apionidae	Coleoptera
<i>Protapion filirostre</i>	Apionidae	Coleoptera
<i>Protapion fulvipes</i>	Apionidae	Coleoptera
<i>Protapion nigritarse</i>	Apionidae	Coleoptera
<i>Diplapion stolidum</i>	Apionidae	Coleoptera
<i>Demetrias atricapillus</i>	Carabidae	Coleoptera
<i>Harpalus affinis</i>	Carabidae	Coleoptera
<i>Licinus punctatulus</i>	Carabidae	Coleoptera
<i>Paradromius linearis</i>	Carabidae	Coleoptera
<i>Altica lythri</i>	Chrysomelidae	Coleoptera
<i>Aphthona euphorbiae</i>	Chrysomelidae	Coleoptera
<i>Bruchidius villosus</i>	Chrysomelidae	Coleoptera
<i>Bruchus rufipes</i>	Chrysomelidae	Coleoptera
<i>Crepidodera aurata</i>	Chrysomelidae	Coleoptera

Species	Family	Order
<i>Crepidodera fulvicornis</i>	Chrysomelidae	Coleoptera
<i>Cryptocephalus moraei</i>	Chrysomelidae	Coleoptera
<i>Cryptocephalus pusillus</i>	Chrysomelidae	Coleoptera
<i>Longitarsus flavicornis</i>	Chrysomelidae	Coleoptera
<i>Longitarsus gracilis</i>	Chrysomelidae	Coleoptera
<i>Longitarsus luridus</i>	Chrysomelidae	Coleoptera
<i>Phyllotreta vittula</i>	Chrysomelidae	Coleoptera
<i>Coccinella septempunctata</i>	Coccinellidae	Coleoptera
<i>Propylea quattuordecimpunctata</i>	Coccinellidae	Coleoptera
<i>Psyllobora vigintiduopunctata</i>	Coccinellidae	Coleoptera
<i>Tytthaspis sedecimpunctata</i>	Coccinellidae	Coleoptera
<i>Halyzia sedecimguttata</i>	Coccinellidae	Coleoptera
<i>Calvia quattuordecimguttata</i>	Coccinellidae	Coleoptera
<i>Adalia decempunctata</i>	Coccinellidae	Coleoptera
<i>Harmonia axyridis</i>	Coccinellidae	Coleoptera
<i>Rhyzobius litura</i>	Coccinellidae	Coleoptera
<i>Cryptophagus dentatus</i>	Cryptophagidae	Coleoptera
<i>Anthonomus rubi</i>	Curculionidae	Coleoptera
<i>Coelositona cambricus</i>	Curculionidae	Coleoptera
<i>Curculio glandium</i>	Curculionidae	Coleoptera
<i>Glocianus punctiger</i>	Curculionidae	Coleoptera
<i>Mecinus pascuorum</i>	Curculionidae	Coleoptera
<i>Sitona hispidulus</i>	Curculionidae	Coleoptera
<i>Sitona lineatus</i>	Curculionidae	Coleoptera
<i>Sitona sulcifrons</i>	Curculionidae	Coleoptera
<i>Trichosirocalus barnevillei</i>	Curculionidae	Coleoptera
<i>Tychius junceus</i>	Curculionidae	Coleoptera
<i>Tychius picirostris</i>	Curculionidae	Coleoptera
<i>Cartodere bifasciata</i>	Latridiidae	Coleoptera
<i>Corticarina minuta</i>	Latridiidae	Coleoptera
<i>Corticara gibbosa</i>	Latridiidae	Coleoptera
<i>Oedemera lurida</i>	Oedemeridae	Coleoptera
<i>Olibrus aeneus</i>	Phalacridae	Coleoptera

Species	Family	Order
<i>Olibrus affinis</i>	Phalacridae	Coleoptera
<i>Cypha longicornis</i>	Staphylinidae	Coleoptera
<i>Metopsia clypeata</i>	Staphylinidae	Coleoptera
<i>Philonthus cognatus</i>	Staphylinidae	Coleoptera
<i>Stenus fulvicornis</i>	Staphylinidae	Coleoptera
<i>Stenus ossium</i>	Staphylinidae	Coleoptera
<i>Tachyporus chrysomelinus</i>	Staphylinidae	Coleoptera
<i>Tachyporus hypnorum</i>	Staphylinidae	Coleoptera
<i>Adia cinerella</i>	Anthomyiidae	Diptera
<i>Anthomyza gracilis</i>	Anthomyzidae	Diptera
<i>Machimus atricapillus</i>	Asilidae	Diptera
<i>Pollenia rudis</i>	Calliphoridae	Diptera
<i>Stevenia deceptor</i>	Calliphoridae	Diptera
<i>Chamaemyia herbarum</i>	Chamaemyiidae	Diptera
<i>Sicus ferrugineus</i>	Conopidae	Diptera
<i>Physocephala rufipes</i>	Conopidae	Diptera
<i>Chrysotus gramineus</i>	Dolichopodidae	Diptera
<i>Dolichopus festus</i>	Dolichopodidae	Diptera
<i>Dolichopus griseipennis</i>	Dolichopodidae	Diptera
<i>Scaptomyza graminum</i>	Drosophilidae	Diptera
<i>Discomyza incurva</i>	Ephydriidae	Diptera
<i>Hydrellia griseola</i>	Ephydriidae	Diptera
<i>Psilopa nitidula</i>	Ephydriidae	Diptera
<i>Trimerina madizans</i>	Ephydriidae	Diptera
<i>Minettia fasciata</i> (Fallén, 1826)	Lauxaniidae	Diptera
<i>Lonchoptera lutea</i>	Lonchopteridae	Diptera
<i>Coenosia infantula</i>	Muscidae	Diptera
<i>Coenosia tigrina</i>	Muscidae	Diptera
<i>Musca autumnalis</i>	Muscidae	Diptera
<i>Muscina levida</i>	Muscidae	Diptera
<i>Orchisia costata</i>	Muscidae	Diptera
<i>Schoenomyza litorella</i>	Muscidae	Diptera
<i>Geomyza balachowskyi</i>	Opomyzidae	Diptera
<i>Opomyza petrei</i>	Opomyzidae	Diptera

Species	Family	Order
<i>Chrysopilus cristatus</i>	Rhagionidae	Diptera
<i>Blaesoxipha plumicornis</i>	Sarcophagidae	Diptera
<i>Ravinia pernix</i>	Sarcophagidae	Diptera
<i>Sarcophaga anaces</i>	Sarcophagidae	Diptera
<i>Sarcophaga nigriventris</i>	Sarcophagidae	Diptera
<i>Limnia unguicornis</i>	Sciomyzidae	Diptera
<i>Pherbellia cinerella</i>	Sciomyzidae	Diptera
<i>Tetanocera elata</i>	Sciomyzidae	Diptera
<i>Sepsis fulgens</i>	Sepsidae	Diptera
<i>Chorisops nagatomii</i>	Stratiomyidae	Diptera
<i>Cheilosia pagana</i>	Syrphidae	Diptera
<i>Cheilosia proxima</i>	Syrphidae	Diptera
<i>Eristalis arbustorum</i>	Syrphidae	Diptera
<i>Eristalis tenax</i>	Syrphidae	Diptera
<i>Eupeodes corollae</i>	Syrphidae	Diptera
<i>Eupeodes luniger</i>	Syrphidae	Diptera
<i>Helophilus pendulus</i>	Syrphidae	Diptera
<i>Neoscia podagrica</i>	Syrphidae	Diptera
<i>Pipizella viduata</i>	Syrphidae	Diptera
<i>Platycheirus albimanus</i>	Syrphidae	Diptera
<i>Platycheirus scutatus</i>	Syrphidae	Diptera
<i>Sphaerophoria scripta</i>	Syrphidae	Diptera
<i>Sphaerophoria taeniata</i>	Syrphidae	Diptera
<i>Syritta pipiens</i>	Syrphidae	Diptera
<i>Syrphus ribesii</i>	Syrphidae	Diptera
<i>Syrphus vitripennis</i>	Syrphidae	Diptera
<i>Aplomya confinis</i>	Tachinidae	Diptera
<i>Ectophasia crassipennis</i>	Tachinidae	Diptera
<i>Eriothrix rufomaculata</i>	Tachinidae	Diptera
<i>Phasia pusilla</i>	Tachinidae	Diptera
<i>Siphona geniculata</i>	Tachinidae	Diptera
<i>Voria ruralis</i>	Tachinidae	Diptera
<i>Sphenella marginata</i>	Tephritidae	Diptera
<i>Tephritis divisa</i>	Tephritidae	Diptera

Species	Family	Order
<i>Terellia longicauda</i>	<i>Tephritidae</i>	<i>Diptera</i>
<i>Herina nigrina</i>	<i>Ulididae</i>	<i>Diptera</i>
<i>Bombus lapidarius</i>	<i>Apidae</i>	<i>Hymenoptera</i>
<i>Bombus pascurom</i>	<i>Apidae</i>	<i>Hymenoptera</i>
<i>Lasius niger</i>	<i>Formicidae</i>	<i>Hymenoptera</i>
<i>Blastobasis rebelii</i>	<i>Blastobasidae</i>	<i>Lepidoptera</i>
<i>Agapeta hamana</i>	<i>Tortricidae</i>	<i>Lepidoptera</i>
<i>Cacoecimorpha pronubana</i>	<i>Tortricidae</i>	<i>Lepidoptera</i>
<i>Grapholita compositella</i>	<i>Tortricidae</i>	<i>Lepidoptera</i>
<i>Pieris brassicae</i>	<i>Pieridae</i>	<i>Lepidoptera</i>
<i>Pieris rapae</i>	<i>Pieridae</i>	<i>Lepidoptera</i>
<i>Polyommatus icarus</i>	<i>Lycaenidae</i>	<i>Lepidoptera</i>
<i>Aricia agestis</i>	<i>Lycaenidae</i>	<i>Lepidoptera</i>
<i>Celastrina argiolus</i>	<i>Lycaenidae</i>	<i>Lepidoptera</i>
<i>Vanessa cardui</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Vanessa atalanta</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Pararge aegeria</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Maniola jurtina</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Pyronia tithonus</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Coenonympha pamphilus</i>	<i>Nymphalidae</i>	<i>Lepidoptera</i>
<i>Pyrausta purpuralis</i>	<i>Crambidae</i>	<i>Lepidoptera</i>
<i>Agriphila straminella</i>	<i>Crambidae</i>	<i>Lepidoptera</i>
<i>Euclidia glyphica</i>	<i>Erebidae</i>	<i>Lepidoptera</i>
<i>Tyria jacobaeae</i>	<i>Erebidae</i>	<i>Lepidoptera</i>
<i>Acronicta rumicis</i>	<i>Noctuidae</i>	<i>Lepidoptera</i>

---

## REFERENCES

<sup>1</sup> Greengage Environmental Ltd (2024) 552979jh06Dec24FV03\_PEA

<sup>2</sup> Red Data Book (RDB) <https://www.iucnredlist.org/>

<sup>3</sup> BSI, (2013). British Standard 42020:2013: Biodiversity – Code of practice for planning and development, BSI Standards Publication.

<sup>4</sup> Chartered Institute of Ecological and Environmental Management (CIEEM) (2019) On The Lifespan Of Ecological Reports & Surveys

<sup>5</sup> Hampshire County Council. (2019); Ecological Appraisal & Phase II Protected Species Surveys.

<sup>6</sup> Ecology & habitat management Ltd. (2023); Preliminary Ecological Assessment, Reptile Survey & Bat Survey Report- Phase 1.

<sup>7</sup> Brooks, S.J. 1993. Joint Committee for the Conservation of British Invertebrates: Guidelines for Invertebrate Surveys. *British Wildlife*, 4(5) 283-287